



The Chain of Chirality Transfer as Determinant of Brain Functional Laterality Breaking the Chirality Silence: Search for New Generation of Biomarkers; Relevance to Neurodegenerative Diseases, Cognitive Psychology, and Nutrition Science

Victor V. Dyakin, PhD^{1*}; Justin Lucas²; Nika V. Dyakina-Fagnano, DO³; Eva V. Posner⁴; Csaba Vadasz, PhD¹

¹The Nathan S. Kline Institute for Psychiatric Research, New York State Office of Mental Health, USA

²Targeted therapeutics Oncology Research Unit, Pfizer, USA

³Child, Adolescent and Young Adult Psychiatry, Columbia University Medical Center, USA

⁴Phelps Hospital, Northwell Health; The Baby-Friendly Hospital Initiative, USA

Abstract

Biomolecules are the products of an evolutionary history. As a result, the phenomenon of molecular chirality is relevant to protein folding, neuronal proliferation, brain functional laterality, as well as the nature of cognition, consciousness, behavior, and psychiatry. Molecular chirality, discovered by Faraday (1846) and Pasteur (1848), helped to reveal that the biochemistry of the living beings has a prevalent chirality. At present, the essence of biochirality is widely recognized, appreciated, and exploited in neuroscience and psychology. In a more general sense, molecular chirality is recognized as the universal "Force of Nature." From the formal geometrical perspective, a chain of chiral bifurcations is the chain of the chirality transfer between the molecular micro-, meso-, and macro-scales. Consequently, the symmetry is considered as a critical issue in the brain information processing. The fundamental laws of information theory reflect the relationship between entropy, symmetry and information. At the cellular level, signal transduction mechanism involves the wave of chiral transformations in the process of protein-protein, protein-phospholipids, and protein-DNA interactions. The symmetry dynamics at the molecular and cellular levels are considered in connection to the laterality of cognitive functions. The abnormal symmetry dynamics viewed as a primary reason of an aggregation of mis-folded proteins in the neurodegenerative diseases and psychiatric disorders. The molecular basis of the "symmetry evolution" in the biological systems is a question of interest. In this short review, we briefly summarize advances in the broad field of biochirality connecting two poles of the phenomena: the atomic orbitals and the brain's cognitive function. Analysis of current results allows introducing the new generation of entangled biomarkers ranging from the molecular chirality to laterality of cognitive and executive functions.

Introduction

The essence of biochirality¹ is widely recognized, appreciated and exploited. Partially it is illustrated by chronologically sequencing citations from the publications in the diverse branches of the science. "Almost every biochemical process occurring in the cell of all living organisms is based on some specific, stereoselective interaction between reacting molecules" [1]. "From a chemical point of view, proteins are by far the most structurally complex and functionally sophisticated molecules known" [2]. "Stereospecificity is one of the hallmarks of enzyme catalysis" [3]. "One of the most dramatic aspects of biological systems involving proton transfer is their high stereoselectivity" [4]. "Chirality plays a fundamental role in the activity of biological molecules and broad classes of chemical reactions" [5]. "Controlled mirror symmetry breaking arising from chemical and physical origin is currently one of the hottest issues in the field of supramolecular chirality" [6]. "Chiral recognition is the fundamental property of many biological molecules" [7]. "Chiral compounds...pose the significant impact on the understanding of the origin of life and all processes that occur in living organisms" [7]. "Cell chirality may be a general property of eukaryotic cells" [8]. "Chirality is one of the ubiquitous phenomena

in biological systems" [9]. "Consistent left-right (LR) asymmetry is a fundamental aspect of the bodyplan across phyla" [10]. "Among the most readily observed topological features in natural structures are chirality, hierarchy, and hierarchy of chirality" [11]. "Chirality is a fundamental property and vital to chemistry, biology, physics and materials science" [12]. In most general sense, molecular chirality is appreciated as the universal "force of nature" [13]. "How chirality at one length scale can be translated to asymmetry at a different scale is largely not well understood" [12].

In such situation, it is reasonable to view the enormous amount of facts of biochirality as the hierarchical system unifying by the common underlying mechanism.

Stereospecific Phenomena

Chirality

The chiral, spiral and helical structures and shapes are seen in the morphology of many physical [15] and biological [16] systems including the spiraling of plants [17] the shape of the mammalian cochlea [18], and the shape of human brain [8]. In any organism, the geometrical patterns are seen in a position of the heart and visceral organs [10], and in the spatial distribution of brain activity [19]. Left-right asymmetry recognized as a fundamental property of the brain, evident at all levels of an organization, including molecular, cellular, morphological, and functional [20-23]. At the same time, it is well known that the majority of biologically active molecules such as amino acids [22], proteins [23], carbohydrates,¹¹ and phospholipids [24] are chiral. The ability of proteins to fold into the chiral sec-

*Correspondence: Victor V. Dyakin, PhD

The Nathan S. Kline Institute for Psychiatric Research, New York State Office of Mental Health, 140 Old Orangeburg Road, Bldg. 35. Orangeburg, New York, USA. Email: dyakin@nki.rfmh.org

Received: May 31, 2017; Accepted: August 17, 2017; Published: September 19, 2017

Neurology and Neuroscience Research. 2017;1(1):2
DOI: [10.24983/scitemed.nnr.2017.00028](https://doi.org/10.24983/scitemed.nnr.2017.00028)

Copyright © 2017 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY).

Neurology and Neuroscience Research. 2017;1(1):2

¹We assume that the readers are familiar with the basic concept of the chirality in relation to the theory of symmetry groups, quantum electrodynamics, and stereochemistry [14].

¹¹Carbohydrates contain multiple stereocenters, allowing many forms of isomers including enantiomers, diastereoisomers, and epimers.

ondary, tertiary, and higher-ordered structures assumed to be responsible for their prominent role in the chirality transformation events. The members of protein family known for chirality related functions include enzymesⁱⁱⁱ [25], cytoskeleton molecular complex [26], amylogenic proteins [27,28], trans-membrane proteins – ligand complex [29], drugs^{iv} [30,31], and antibiotics^v [32,33]. For the purpose of our review we will restrict our attention mostly to the proteins and phospholipids.

Chirality Prevalence (Amino Acids, Sugars, Phospholipids, Water)

Most of the bio-compounds have prevalent chirality (phenomenon of homochirality).^{vi}

AMINO ACIDS/PROTEINS. Among more than 700 naturally occurring amino acids only 20 are involved in the proteins synthesis. All of the alpha-amino acids are chiral (except glycine). The essential chiral amino acids are L-configurations^{vii} ("left-handed"). The amino acids stereo-chemistry is the major determinant of protein chirality. It is meaningful that the spatial configuration of amino acids in the organic world, as a rule, is governed by enzymes (i.e. proteins) [25]. The evolution known plays a pivotal role in the mechanism of protein folding. Chiral proteins, as the principal constituents of neuronal cells, provide molecular machinery for the lateralized brain cognitive functions.

SUGARS. The sugars are D-configurations ("right-handed").

PHOSPHOLIPIDS. The majority of the membrane phospholipids^{viii} is right-handed, but in the archaea (single-celled organisms) membrane they are left-handed [24,34]. Furthermore, the membrane phospholipids are right-handed [24,34] but in an archaea (single-celled organism)^{ix} membrane, they are left-handed.

WATER. The water molecules (H₂O) as the main constituent of the living organism are in the scope of our consideration. In a homogeneous, achiral environment, the H₂O molecules possess neither a chiral center nor a helical conformation that can cause spontaneous chirality effects. Accordingly, many assume (at first glance) that water falls apart from phenomenon bio-chirality. However, this view was challenged by studies that devote more close attention to the dynamics of water structure in the chiral environment (proteins, phospholipids, polysaccharides) and dynamics of chirality. In the variety of the "inhomogeneous" situations, water molecules dynamically participate in chirality-related effects. Among such conditions, we can point on the aggregation of water molecules into clusters [38], interaction with magnetic bio-proteins [39], contact with the solid/gel/liquid surface [40], interaction with the chiral solutes [41,42], and dehydration synthesis of proteins (via covalent bonds of amino-acids). In particular, it was shown that structured water exhibits a chirality adapted from DNA [43]. The presence of hydrophobic, hydrophilic, and intermediate groups in the amino acids and phospholipids contributes to the variety pathways of spatial arrangement. The body of accumulated evidence (despite the variety of distinct pathways) suggests some common causal agent in the chain of events from the molecular chirality to the origin of body/brain morphology and function. This "universal agent" is the spatial relationships between the objects linked to the fundamental geometrical patterns of space and force fields.

Biological Evolution as a Chain of Chiral Bifurcations

The pathway of life can be perceived as an "increasing chemical and physical complexity" [44,45]. In a more particular sense, the biological evolution considered as the chain of chiral bifurcations [46].

The symmetry effects propagate from the level of molecular spatial

transformations, intracellular /cross-membrane molecular transport, cell motility, to the higher hierarchical level of the spatial organization such as the cell proliferation [47], immune defense, environmental chemistry [49], motor behavior, brain cognitive functions, human psychology [50], psychiatry [48,51-53]. The symmetry effects play a critical role in food preparation [54,55], pharmaceutical industry [56,57], design of molecular devices (biosensors and information processing units). Symmetry is considered the critical issue in the theory of information processing and the design of artificial intellect^x [58]. The hierarchy of biological structural organization arms an organism with the abilities of adequate response to environmental challenges of differential time-space ranks.

Origin of Biological Homochirality

The homochirality is usually associated with bio-molecular objects (distinct from non-biological objects). There is an objective reason for this specificity. The biological homochirality known to be essential for the molecular recognition, protein replication, post-translational modification, and degradation processes.

The origin of homochirality in biology is the subject of much debate. The emotional view on bio-chirality is frequently associated with an expression such as a symmetry breaking,^{xi} enigma, puzzle, or mystery. More rational scientific approaches exhibit some degree of uncertainty, but always are ready to move forward along with the newly-coming objective evidence. After works of Prelog, the discussion of an origin and maintenance of homochirality was shifted from the intuitive, emotional domain to the ground of scientific facts [59]. The cumulative advance in chiral physics, chemistry, and mathematics sheds light on the phenomenon of chirality in general and on biological homochirality in particular. Several decades ago Frank developed the mathematical model^{xii} for the spontaneous autocatalytic reaction as the mechanism for the evolution of homochirality [60,61]. Recently it was shown that the origin of molecular homochirality could be attributed to the non-equilibrium state of biological systems. The Vester-Ulbricht hypothesis, based on the interaction of the left-handed electrons (present in beta-radiation) with biological materials which preferentially destroyed one of the two enantiomer, was the first step in the right direction [62,63]. This hypothesis was successfully tested experimentally [64]. The sensitivity of molecular chirality to external determinants becomes one of the most productive ideas. The classic experiment of Viedma reveals that the bulk and dispersed solid state crystals show different dynamics of solid-liquid phase transition, which, under certain conditions, can be utilized to achieve an enantiomeric preference [65]. The following studies of the chirality dynamics in the solid-liquid phase system leads to several models presumably providing the mechanism implicating for the origin of biochirality. Among them is the model based on the different solubility of homochiral and heterochiral crystals [66] and "chiral amnesia" model [67,68]. The review of current hypothesizes on the biological homochirality in connection with the recent discovery of the interstellar chiral molecule can be found in multiple sources [10,44,69-71]. The new finding reveals the fundamental significance of the symmetry transfer (1) and symmetry-associated phase transitions (2) in resolving the origin of biological homochirality [44,70,72]. Among them are the chiral recognition/selection during the self-assembly of protein-mimic macro-anions [73], chiral recognition in the transmitter-receptor interactions [74], the chirality-induced conformation of the cell membrane lipid rafts^{xiii} [75,76], and the phase-transfer chiral catalysis^{xiv} [77]. The chiral catalysis is productively utilized for symmetry-asymmetry

ⁱⁱⁱ The enzymes determine the crystallization of chiral amino acids [25] and the "handedness" (chirality) of the proteins during catalytic synthesis.

^{iv} The enantiomers of a drug, as a rule, are differing in potency, toxicity, and behavior in biological systems.

^v The advance in enantio-separation of the antibiotics reveals the significance of chirality in their biological activity [32,33].

^{vi} Phospholipids and cholesterol both contain chiral carbon atoms and could themselves mediate stereoselective effects [35].

^{vii} Brain tightly regulates the balance between the levels of right- and left-handed amino acids for proper structure of functional proteins [22,36,37].

^{viii} It is essential to know that homochirality is always relative (not absolute). There are many examples in nature where polypeptide topologies use both: L- and D-amino acids [36,37]

^{ix} Archaea are classified as microbes (single-celled prokaryotes).

^x It is notable that the design of an intelligent human-made sensing system is based on the discrimination of the object chirality and the use of chiral shape-defined polymers such as the dynamic helical polymers [58].

^{xi} At the molecular level, the term "symmetry breaking" refers to the imbalance between two enantiomers.

^{xii} The main idea is that a substance acts as a catalyst in its own self-production and at the same time acts to suppress synthesis of its enantiomer.

^{xiii} It has been suggested that the chiral nature of cholesterol plays a role in the process of the bud formation [85].

^{xiv} The variety of intracellular chemical reactions is mediated by seospecific enzymes. Many catalytic mechanisms require stereospecific deprotonation and reprotonation steps [77]. The L-amino acid active center is the primary determinant of catalytic activity. Thus, L-amino acid actively transfer chirality to the stereospecific enzyme activities [77].

transformations in biochemistry [78]. The phenomenon of chirality is common to all aspects of animal life and body/brain morphology, including the internal organs, the sensory systems, the central nervous system, and behavior. In particular, it pertains to sensory perception [79], motor-behavior [80], and food consumption [54]. All of the arguments related to the origin of the biological homochirality, including terrestrial [44] and extraterrestrial [81,82] are based on the common and experimentally proved effects. The sensitivity of chiral molecules to the internal, external, and mutual (internal-external) physical (objective) parameters [83]. While you can find discussions of the many particular local physical parameters, in the majority of current reviews, you will almost never find discussions about of the global parameters, such as the chirality of space-time evident in the physical systems [83,84].

Transformation of Bio-Chirality

Symmetry transformation is traditionally considered to be the spontaneous processes accompanying all of the molecular interactions (phase separation and phase transition), including symmetry breaking and molecular folding in relation to information minimization or symmetry maximization [86]. The symmetry dynamics at the molecular level are frequently considered to be the mechanism responsible for the laterality of cognitive functions. Thus, the laterality and hemispheric asymmetry should be regarded as the indispensable attributes of the brain. The molecular basis of the symmetry evolution in biological systems is the question of interest. Current evidence suggests that integration of the internal and external determinants [84], as well as global and local signaling pathways, is necessary for orienting the diverse levels of structure with respect to the body/brain axes [87-89]. In particular, the integration is evident in the hierarchical chirality transformation accompanying the planar cell polarity^{xv} in the epithelium, sensory organs [90],^{xvi} and the brain morphology [89].

Dynamic Organic Reactions

The spatial interpretation of molecular structures was advanced (1874) by the concepts of the tetrahedral orientation of carbon's four bonds by Van't Hoff and optical isomerism by Le Bel [91,92]. These ideas, in turn, gave rise to the stereochemistry of chiral stereoisomers. It was realized that the spatial orientation of different functional groups governs the patterns of specificity in chemical reactivity [93]. The interaction between the chiral and achiral components was widely studied in the biological organic reactions. The dynamics of organic reactions (in particular those that take place within the spatially ordered environment of an enzyme protein interaction) exhibit stereoselectivity and stereospecificity. The excellent review of the stereo-dynamic of chiral objects, including natural molecular structures and artificial molecular devices, can be found in [94]. It is notable, that both the stereoselectivity and stereospecificity are based on the recognition of the symmetry-related characters (such as polarity, chirality, and helicity). Thus, a conclusion such as "chiral recognition is the fundamental property of many biological molecules" is not surprising [7].

Diversity of Stereospecific Phenomena

The diversity of stereospecific (chiral) phenomena was observed in molecular structures including chirality recognition/sensing [95,96], chirality transfer [97], chirality/helicity induction [98-101], chirality amplification [102-104], chirality breaking [61,105], chirality conflict [106,107], helicity inversion [108-110], and chiral phase transitions^{xvii} [111-114,116]. Progress in the studies of molecular chirality transformation is helpful in resolving three questions. First, what kind of determinants can provide a favor in the production of one enantiomer over the other? Second, what is the mechanism of intermolecular propagation and preservation of chirality? And third, what is the mechanism of chirality propagation from the molecular level to a higher degree of biological organization?

Chiral Phase Transitions in Relation to Chirality Transfer

Physical System

Among the variety of the topological phase transitions [120-123] several sub-categories, including the order-disorder [118], chirality-related [119] and geometry-induced [120,123] transitions, were discovered. In condensed matter physics, phase transitions exhibit sensitivity to external physical parameters (pressure, fields, electro-magnetic radiation, sonication,^{xviii} and doping). The systematic study of these phenomena has revealed a new class of phase transitions, called "quantum phase transition (QFT)". Spontaneous QFTs take place at the mesoscopic level at zero temperature and are driven by the quantum fluctuations^{xix} (according to Heisenberg's uncertainty principle). For the two above facts, the sensitivity to external physical parameters and the concept of the spontaneous phase transitions are of great importance.

Biological Systems

In relation to the biological systems, the chirality of the sub-cellular structures (such as hair bundles) was observed in auditory and vestibular sensory neurons of vertebrates [87,118]. The phase transitions are the common effects, observed in the chiral molecules with pyramidal atomic centers such carbon and nitrogen. The frequency of a pyramidal inversion (tunnel quantum-mechanical effect) depends on the value of the energy barrier and set of external physicochemical factors. Chirality transfer from molecular to morphological level is observable in a diversity of physical objects as well as biochemical and synthetic materials [124,125]. The hierarchical propagation of chirality was found between the objects of different size, shape, and dimensionality [126]. The chirality breaking in the nonequilibrium systems (Bloch walls) was studied in magnetic materials [127]. Recent progress in the development the concept of "active fluids" reveals the chiral behavior of a class of nonequilibrium systems, which include bacterial suspensions of a bacteria, cytoskeleton proteins, and biological tissues. Even relatively simple combination of chiral and achiral stresses, leads to an "unprecedented range of complex motilities, including oscillatory swimming, helical swimming, and run-and-tumble motion" [128].

Chirality Transfer

Variety of Spatially Related Events

Among the variety of spatially related events, the phase transitions and the symmetry transformations are known as the most closely associated. We will focus mostly on the phenomena of the chirality transfer. The chirality transfer from the molecular to morphological scales was documented in nano-materials [124] and polymers [125]. The studies of the plant's growth reveal the "hierarchy of chirality" which transfers from the molecular (lower) levels to the macro-morphological (higher) level [130]. In our review, we explore what is currently known about how the molecular chirality is transformed into to the laterality of cognitive functions.

Sensitivity of Chirality Transfer to Internal, External, Local and Global Conditions

Symmetry transformation arising from chemical and physical origin is one of the hottest issues in the field of molecular chirality [5-8]. Before the further consideration it is essential to emphasize that hypothesis about the distinction between internal and external determinates in the living organism being very productive could become irrational in the form of an absolute opposition. The most convincing example of the link between external and internal factors is the function of a digestive system, which permanently transforms the external factors into the very internal. Pasteur (1861-1887) was the first who introduced an idea regarding an interaction between the polar physical fields (electrical, magnetic, electromagnetic, gravitational/mechanical) with atomic/molecular chirality [131]. He

^{xv} Planar cell polarity (PCP) "is driven by multiple global cues, including gradients of gene expression, gradients of secreted ligands, and anisotropic tissue strain" [117].

^{xvi} The chirality of the sub-cellular structures (such as hair bundles) was observed in auditory and vestibular sensory neurons of vertebrates [87,118].

^{xvii} The formation of supramolecular helicity is associated with the occurrence of new thermodynamic phases. The chirality related effects in thermotropic and lyotropic liquid crystals were demonstrated through the ferroelectricity [115]. The phase transitions were shown to be implicated in the effects of the chirality transformation and monochirality observed in the amino acids and proteins [119].

^{xviii} The sonication-induced chiral symmetry breaking events were observed during sol-gel phase transition [129].

^{xix} The quantum fluctuations are associated with Heisenberg's uncertainty principle.

was also the first who realize the fact of the chirality transfer from the molecular level to the level of macroscopic solid crystal [132]. The ideas were great, but the success was limited. Much later the empirical evidence of the role of the forces of nature on the chiral compounds was obtained. The early idea of Pasteur was supported by modern experimental capabilities [133,134]. Two essential facts were observed: the effect of magnetic field on the molecular chirality [135] and increased enzyme stereo specificity in the course of an enzyme/protein evolution [136,137]. Both results, accompanied by the discovery of the spontaneous [138] and induced chirality provided the tools to disclose the previously mysterious homochirality of life [139]. In particular an essential result was derived by Barron [140]. He demonstrated that supramolecular helices formed from achiral monomers have been controlled by applying the combination of the gravitational and rotational forces [140]. The transfer of stereochemical information (in the form of chirality and helicity) was observed between chiral and achiral constituents of the molecular complexes in general and of biological systems in particular [141-144]. The transfer of chirality from protein to the cellular and embryonic level was suggested in several studies [145,146]. The critical role of a molecular and cellular chirality as the determinants of LR asymmetric in the animal body, and functions has gradually emerged [8]. The explosive advance in the study of asymmetric catalysis over the last four decades has dramatically altered the view on the biomolecular chirality dynamics [147,148]. Before the era of chiral catalysis, the most common characteristic of the enantiomers was the absence or little difference in the chemical and physical properties. At present at the majority of publications related to chiral catalysis we can meet the statements like "enantiomers have different properties". Recent progress in the synthesis of the chiral compound associated with the study of the catalytic asymmetric reactions of carbonyl compounds, allow an understanding of the principles governing the dynamics of structural conformations in the amino acids and proteins of the living organisms [149]. Molecular chirality and correspondently a chirality transfer are recognized as sensitive to the broad range of modulators including the internal, external, localized, and diffused determinants. It is notable that molecular chirality exhibits sensitivity to all types of the chemical binding including ionic, covalent, and not-covalent [58].

The generation of chiral imbalance in the chiral molecular systems can occur spontaneously, due to intrinsic instability or induced by external factors [102]. In accordance with this instability, the chiral self-organization of molecular complexes is sensitive to the impact of many external factors including electrical (metal ions), magnetic, electromagnetic (photon), mechanical, and gravity force fields. The discovery of magnetically induced optical activity by Faraday was the first demonstration of the sensitivity of a molecular chirality to the physical parameters of an environment^{xx} [150]. Since then, the sensitivity of the chiral objects to the environmental parameters has been explored at cosmological [151], molecular [152], atomic [57], and elementary particles levels [153]. Thus, it is reasonable to be aware that different molecular structure can have the same or different physical properties depending on the nature of the physical effect and chemical environment.^{xxi} In the specific case of the stereoisomers interaction with an electromagnetic field, we have at last three different situations depending on the energy diapason (such as IR, UV and NMR spectra), and method used (such as the circular dichroism) [40]. The photons (chiral object itself) of different energy interact with the chiral components of the molecular complex (such as electron or proton) compromising its equilibrium spatial configuration. The recent discovery of the quantum chiral light-matter interaction offers fundamentally new functionalities for the chirality transfer of the bio-molecular structure related to brain quantum information-processing capability [160]. The chiral molecules reveal the capability of the self-organization of the helical

superstructures. The intermolecular interactions related to the modulation of chirality are the part of the supramolecular chemistry [161-165] and interfacial sciences [166,167]. The chiral sensing based on the concept of chirality transfer is of great importance.

Chain of Chirality Transfer

Several relatively new fields of science provide the bridge between dynamic chirality in solid matter physics and bio-chirality. The chirality transfer (or the transfer of handedness) is observed between organic and inorganic molecular structure [168]. The central point of these studies is the chirality transfer in the variety of forms.^{xxii} Among them, we can mention the stereo-physics of liquid crystals [126] and chiral catalysis [169]. The modeling macroscopic chirality emerged from the chiral molecular elements is a challenge for theory, computations, and experiments [126]. Numerous experimental results demonstrate the transfer of chirality among different length scales ranging from dimensions of the elementary particles to the macro-scale (the length of the axon) [124]. In particular, it was shown that the chirality at the molecular scale (amino, acids, proteins, and polysaccharides) could be transferred to the macroscopic and macro-level (neurofilaments and inorganic crystals) as shown in [168]. The issue of dimensionality in the chirality transfer effects is critical for brain information processing in the brain and artificial intelligence devices.^{xxiii} The examples of macroscopic chirality are found in the plant kingdom, animal kingdom and all other groups of organisms.

Physical Systems

In the quantum spin systems, the symmetry-related phase transitions [78,154,155] and the transfer of the stereospecific (symmetry) characters^{xxiv} [97,156] are well-known phenomena. The recent advance in the experimental and theoretical areas of many disciplines related to stereochemistry revealed the chirality-induction effects in the various inorganic materials with mono-chiral and hybrid-chirality structures including plasmonic, semiconducting, metal oxide and silica-based compounds [170]. The most prominent among the field-induced chirality effects are the following: Coulomb (near-field, dipolar), electromagnetic, and plasmonic mechanisms [171]. The chirality transfer from the spin-quantum system of elementary particles to the atomic structure level is an essential element of basic knowledge and serves as the necessary introduction to the understanding of the chemistry and biochemistry. The nuclei of atoms and associated electron system have an innate chirality. The chemical phenomena are viewed as associated with the chirality of electron system and nuclear^{xxv} constituents [172]. The molecular chirality is the consequence of the chirality transfer from the dynamic complex of elementary particles. The transfer of molecular chirality from monomers to polymeric structures has been widely explored and utilized [173]. The advanced field of chiral photon-electron/proton interaction reveals the sensitivity of chiral objects to the physical environment. As a result of such sensitivity, chiral photonelectron/proton interaction offers fundamentally new functional opportunities for information transfer technology and info-processing systems. In particular, the non-reciprocal single-photon devices allow utilizing the quantum information processing based on the superposition of two operational states in chiral spin-photon system [160]. Thus, the stereo-specific effects, including the chirality transfer, are not the unique properties of the organic world.^{xxvi} Quite contrary stereospecific effects are the universal and fundamental character of both organic and inorganic materials.

Biological Systems: Basic Set of the Chirality Transfer Levels

From the physical world chiral events diversity and complexity point of view our primary concern is the chain of the chirality transfer in biological

^{xx} Following Faraday's discovery [150], Pasteur tried to grow chiral crystals in the presence of magnetic field [157].

^{xxi} The NMR spectra of two stereoisomers can be identical or differ depending on the chirality of an environment (solvent).

^{xxii} The effect of chirality transfer is routinely used in nuclear magnetic resonance spectroscopy for the discrimination of the molecular stereoisomers absolute configurations [158].

^{xxiii} The chirality transfer from monomers to a polymer is widely used in plastic engineering [159].

^{xxiv} The process is frequently referred to as the chirality transfer.

^{xxv} "All nuclei are innately chiral and, because electrons can penetrate nuclei, all atoms and molecules are likewise chiral" [172].

^{xxvi} It is notable for the sake of our review that the spiral galaxies are the objects showing preferred chirality analogous to amino acids and sugars [175]. According to a contemporary view, the origins of the elementary particles and chemical elements are associated with the physical events at the galactic scale [176].

systems. The very essential prediction of the sequential chain of chiral events in the organism was done long before the modern progress in biostereochemistry [174]. Taking the review of newly discovered facts as a basis, we will clarify the natures of elements in this chain and the hierarchy of these elements within the chain. Referring to the hierarchy of a chirality transfer, we will assume (based on the review of current publications) that it consists of several distinct levels. The basic set of these levels includes the transfer of following types:

- From the elementary particles to the atomic orbital level
- From the atomic orbital to the molecular level^{xxvii}
- From molecular to macro- and supra-molecular level
- From the molecular level to the cellular level
- From the cellular level to neuronal circuits level
- From the cellular to morphological level
- From the morphological to cognitive level
- From the cognitive to the behavioral level

After reviewing the elements of the chiral hierarchy, we will examine what is currently known about the sensitivity of each of the hierarchical levels of chirality to the internal and external determinants. The chirality transfer occurs under the influence of physical and chemical determinants which play a role of the “chirality directing force.” The stochastic fluctuations in parameters of an environment could result in the transient fluctuations in the relative prevalence of enantiomers [178], while long-term fluctuations will lead to permanent effects giving a chance for amplification and preservation mechanisms. Several reviews provide information about the range of chirality related events [44,45]. Finally, the degree of “stereo-sensitivity” will reveal the multi-variable pathway contributing to the evolution of the brain cognitive functions.

From Elementary Particles to Atomic Level

The chirality transfer from the elementary particles to the arrangement of atomic orbital was considered in the previous paragraphs. The energy difference (parity violation) between the ground and excited states of molecular enantiomers in the presence of weak nuclear force is predicted by theory and proved experimentally [179,180].

From Atomic Orbitals to Molecular Level

The wavefunctions of electron orbitals are traditionally considered to be the determinants of the molecular chirality [14]. Consequently, in the stereochemistry, the spatial arrangement of the atomic orbital is the primary determinant of the chiral center's function in bio-molecules, including the amino acids, sugars, and phospholipids. At present, it is a common recognition that the electronic orbitals of the carbon atom constitute the root contributing to the molecular chirality^{xxviii} [181,182].

From Molecular to Macro-Molecular and Supra-molecular Level

The alterations of brain molecular chirality, represented in particular by the proteins or lipids constituents, are accompanied by the changes in the left/right asymmetry of the synapse (cellular chirality), asymmetry in the regional brain morphology, and laterality of brain functions in many experimental situations. In this sense the molecular chirality, is the principal initiator of the origin of the life. We will review the chirality transfer events in the order of their natural sequence. The transfer of the symmetry patterns (chirality-induced helicity or chirality-helicity transfer) from the amino acids to peptides [179] and proteins [181] is broadly studied. Four main categories of biological-macromolecules, which exhibit chirality, are

proteins, lipids, carbohydrates, and nucleic acids [180,183].

PROTEINS. In the human body about 100,000 different proteins introduce the chirality phenomenon for all the key physiological, perceptual, cognitive and psychological function of an organism. The chirality transfer from amino acids to proteins secondary and higher order structure is one of the most studied fields in biochemistry. The chirality of protein folding gained attention in condensed matter physics [14,184] and molecular biology [36,185]. The stereo-transformations of proteins are a highly dynamic field of science involving the most advanced analytical capabilities [146,186-196]. The protein's stereo-transformations is the area of particular interest in the neuroscience due to relevance to proteins aggregation disorders such as Alzheimer's, Parkinson's, and Huntington's disease (AD, PD, HD).

CELL MEMBRANE (PHOSPHOLIPIDS AND CHOLESTEROL). A similar mechanism is responsible for the transfer of molecular-level stereo-specificity (chirality) to the supra-molecular level (helicity) in cell membrane rafts during endocytosis [197]. The establishment of helical handedness can be formed at the macro-molecular level due to the stereo-ordering regularity of constituted chiral entities at the intra-molecular and inter-molecular interactions [198]. The chirality transfer from the molecular to the supramolecular level (nanometer and micrometer scale) and the morphological level was observed in the inorganic liquid crystals [115], polymers [198], and bio-polymers^{xxix} including cellulose, sugars, proteins, RNA, and DNA.

From Molecular to Cellular Level

The origin of cell chirality and its role in the upstream laterality are the subject of many reviews [7]. The review of current studies suggests that “molecular chirality direct whole-cell chirality [8]. The chirality transfer from the molecular (cytoskeleton proteins) to the cellular level (cell wall and cell shape) was demonstrated in the bacterium [199,200,201, 202]. Cellular chirality, in a variety of its forms, is governed by crosstalk of the internal and external determinants.^{xxx} Chirality at the cellular level was first studied mostly in ciliates or single-celled protozoans. These studies reveal that molecular chirality directs whole-cell chirality^{xxxi} [8]. At the cellular level, signal transduction involves the wave of chiral transformations in the protein-phospholipid interactions [204-206].

From Cellular to Morphological Level

The chirality of the biomolecules and the intrinsic cell chirality observed in various organisms appear to be a causal event for the left-right (LR) asymmetric of morphogenesis [8,130]. The multiple pathways of chirality transfer from the molecular cytoskeleton dynamics to the cellular behavior, and to organ asymmetry were found [10]. As was mentioned above, the cell-shape chirality (cell chirality) is found to be driven by an intrinsic molecular mechanism comprising from the family of chiral cytoskeleton proteins [203]. In turn, the chirality transfer from the “chiral cells” to an organ occurs by the process of “planar cell chirality” (PCC). According to the early observation of Brown and Wolpert “the spiral asymmetry, as seen in spiral cleavage and ciliates, involves the conversion of molecular asymmetry to the cellular and multicellular level” [174]. In particular, it includes climbing plants tendrils [207,208], flower petals [209], and shells [210]. In the literature, the cellular and morphological hierarchical levels of chirality transfer are frequently considered together. It is notable, that chiral morphology exhibits the prevalence of the handedness [211]. The coherent and quasi-independent role of the intracellular (gene expression and cytoskeleton dynamics) and extracellular (cilia) determinants in

^{xxvii} Due to the topic of our review, we discussed very briefly (just mentioned) the two polar elements in the chain of the chirality transfer: the physics of elementary particles and space-time chirality of an environment. Actually these two elements are seemingly-polar only within our “hierarchical chain”. In the nature they are inherently linked to each other by concept of space-time symmetry. Thus the linear sequence of chiral events, in this consideration, becomes a natural circle of the chiral transformations. For those who are interested in the quantum aspects of the chirality transfer we recommend to explore the “chiral effective field theory” [177].

^{xxviii} Molecular level of the chirality transfer cover the broad range of molecular structures from small molecules (such the amino acids) to the large complexes including polymers (such as proteins) and globular molecules.

^{xxix} Among the biopolymers, we will mostly concentrate on the proteins.

^{xxx} Support of this statement speaks to the fact that the cells isolated from developing organism undergo the symmetry transformation even in the absence of the external signals [8].

^{xxxi} Several chiral compounds exhibit capability to block cell proliferation suggesting their relation to issue of the cancer treatment [47].

the initiation of the embryonic asymmetry were studied in *Caenorhabditis elegans*, *Xenopus*, snails, *Drosophila*,^{xxxii} frogs, chicks, and mice [8,212]. The interaction of proteins and phospholipids chirality at cellular level finally result in the asymmetrical distribution of major CNS receptors^{xxxiii} between left and right cerebral cortex [213].

From the Morphological to Cognitive Level

The chiral, spiral or helical shapes are seen in the morphology of many biological systems, including the spiraling of plants, and shape of the mammalian cochlea [18]. During animal development, chirality transfer repeatedly occurs at the different levels and scales of spatial organization. As we will see later, each lower-level (in the hierarchy of symmetry transformation) generates the higher level of the morphological and functional specialization. The functional specialization of the nervous system at the molecular (intracellular) level is always under impact of the extracellular events. The functions on the cellular and organ (brain) level, in turn, are always under the acute influence of evolutionary preserved behavioral paradigms. The studies of the association of brain function with the body's reflexive reactions and goal-oriented movement were initiated by works of Pavlov in relation to the conditioned and unconditioned reflexes [214]. The ideas of Pavlov resonate with the earliest thoughts of Plate and Aristotle regarding the succession of memory events and modern theories of cognitive association. Brain asymmetry at the structural level begins to be apparent in the fetal brain in humans and nonhuman primates [215]. Brain laterality at the functional level is considered to be a result of morphological laterality [216]. The higher-order cognitive functions are associated with the spatially distributed activity of cerebral cortex. The anatomical asymmetry of the human cerebral cortex is exhibiting a three-partial differentiation. First-the torque {right frontal and left occipital areas are more prominent in size} [217]. Second-the leftward volumetric dominance in language-related areas [218,219]. Third-the left-right asymmetry in cortical thickness (right biased in frontal and left biased in parietal regions [220-222].

The discrimination between sleep and evoke brain activity with fMRI and EEG techniques demonstrates objective (and now trivial) link of the quantum dynamics of proton and electron with the faculties of space perception,^{xxxiv} and consciousness [223,224]. The chirality transfer from the morphological level to functional brain laterality is considered in the plenty of the articles, reviews, and monographs [44,224-226].

From Cognitive to Behavioral Level

Chiral information is used for social communication in the variety of species from insect^{xxxv} to human^{xxxvi} {face expression and perception} [228,229]. Diverse explanations have been proposed for the origin of the behavioral laterality including space constraints [230], as well as genetic, and ecological determinants. The link between the molecular, functional, and behavioral laterality is the focus of many studies. It is known that lateralized expression of the neuro-receptors serves as internal positional markers (compass) to distinguish the left and right hemisphere. Long time ago these markers were hypothesized to be necessary during ontogenesis for bilaterally-symmetrical brain formation and performance the hemisphere-specific perceptual and cognitive functions [231]. In particular, it was shown that the circle rotation of animal, which is mediated by several neurotransmitting systems, significantly contributed by an asymmetric expression of the hypothalamic neurohormones (such as somatostatin, substance P, and others [232]. Thus behavioral studies support the idea of a hierarchical chain of the chirality transfer.

Biological Transport, Information Processing, and Cognitive Systems

It is reasonable to mention here that the chirality transfer is relevant to three essential issues associated with space topology: the biological transport, information processing, and principle of cognition. A wide range of the coordinated motions, including the locomotion of organisms, spatial displacements at morphological (muscle fibers), cellular (flagella, cilia, synaptic spine) and molecular levels (protein fold, lipid chirality) are based on the principles of the chiral dynamics [91]. It has become clear that the evolutionary design of biological molecular motors acts upon the chirality transfer from the amino acids to the helical chirality of proteins. The biological molecular transport systems consist of two major parts: the cytoskeleton and motor proteins [233,234]. The functions of intracellular motor-proteins inspired the engineering molecular motors [235,236]. The advance in the design of artificial molecular motors promotes the understanding of the mechanism of intracellular transport [112,202,237, 238]. Thus not only the lower level biological features like the molecular transport, neuronal signaling, but also the higher level features such as sensory perception, brain information processing, and cognitive functions in general are related to the modality of chirality transfers.

Conclusion

Noether's Theorem

Noether's theorem states that each conservation law^{xxxvii} is associated with symmetry in the underlying physics [239]. The universal role of symmetry principles in the stereochemical configuration traditionally discussed in relations to the fundamental theorems of Noether, Ruch, and Gödel (240,241). From the platform of modern science, each of the conservation and violation laws is fundamentally associated with the preservation or alteration of the symmetry-related parameters. These universal laws are not only relevant to both nonbiological and biological objects but also are the fundamental determinants of an interaction between them. The symmetry patterns are persistently transferred between any combinations of the objects of similar or different nature.

Our Hypothesis

Utilizing the application of universal space time topology principles to the biological world, our hypothesis^{xxxviii} state that the link between molecular chirality and laterality of the cognitive function is based on the universal law of symmetry. Refraining known sentence "space and time are the primary forms of existence" we can say that the dynamic symmetry is the form of existence for the entire variety of the biological objects.

Biological molecules reveal the chirality (handedness) in a variety of fashions many of which are critical for the healthy functions of human body, brain, and mind. The observation of the sign-alternating hierarchies for DNA and proteins contribute to the structure-function link in biochemistry, neuroscience, and psychology [242]. The striking similarity in the chirality-related effects within a broad class of molecular, cellular, and morphological structures inspired researchers in many fields of stereochemistry and neuroscience. The advance in the current results reveals the fundamental significance of molecular chirality in the broad range of inter-related disciplines including an intracellular transport, neuro-development, sensory perception, motor behavior, brain cognitive functions, human psychology, and psychiatry. The helical structures of the proteins are known to be the consequence of amino acid chirality. The helicity of pro-

^{xxxii} In *Drosophila*, intrinsic cell chirality is observed in various left-right (LR) asymmetric tissues, and appears to be responsible for their LR asymmetric morphogenesis. It is notable that the cell chirality can drive the LR asymmetric development of individual organs without establishing the LR axis of the whole embryo [8].

^{xxxiii} The muscarinic acetylcholine receptors (mAChR), the prevalent receptors in the central nervous system, are asymmetrically distributed between left and right cerebral cortex with the right-side dominance [213].

^{xxxiv} In Kant's view, space and time are a priori forms of sensibility, providing two windows' for empirical intuition.

^{xxxv} The insects and many other animals use molecular chirality for olfactory perception. Insects, which use chiral pheromones, typically produce and respond to either a single stereoisomer or to species-specific blend of only some of the possible stereoisomers [227].

^{xxxvi} The face expression and perception, which play a significant role in human social communication, both are the chirality related functions.

^{xxxvii} Conservation Laws: Conservation of the Linear Momentum, Angular Momentum, and Energy.

^{xxxviii} Meaning that the hypothesis is based on the universal nature of Noether's theorem and is generated by the review of contemporary experimental facts and theoretical models.

teins, in turn, provides an opportunity for chiral biological macromolecular systems with higher hierarchical levels of symmetry and yet unknown functions [46,242,243]. The hierarchy in the chain of the chirality transfer, evident particularly from the current review, possesses several not trivial features. Each of the consequently following hierarchical levels is contributed to from the cumulative power of all lower and higher levels. Each level has its own quasi-independent internal and external determinants. The majority of the external determinants (including environmental) exhibit a chiral nature. As such, the chirality of external determinants is a critical factor in the evolution and development of the organism. The chirality is naturally imposed to the brain structure and function from the two seemingly opposite but fundamentally linked to each other sides: the nature of bio-molecular structures and from the geometry of the environment. The fundamental nature of the chirality transfer should be considered in the experimental design strategy in neuroscience, cognitive science, psychology, and pharmacology. In particular, the molecular chirality is a sensitive indicator of both: the proper protein folding and pathological aggregation. Accordingly, the chirality patterns (raging from molecular chirality to the laterality of brain function) should be considered as a new generation of the biomarkers in the spectrum of the disease conditions including neurodegenerative and psychiatric disorders [22,244]

Several recently published essential papers provide the broad review of the previous and current hypothesis of a perceptual and cognitive development in health [245], and pathology [246]. Referring to the limited progress in the current models, and the low efficiency of drug treatment, authors frequently appeal to the necessity of a new approach in the research of the cognitive functions mechanism. Notably that none of the above-mentioned review are focused on the issues of stereochemistry, and brain asymmetry which are a central point for the study of links between protein folding and brain functions. [247-249]. Even the reviews devoted to structure, function, and assembly of the visual system (which is lateralized and well-studied as one of the most asymmetrical) do not focused on the molecular substrate of sensory functions. The explosive attention to protein-brain topology demonstrated in this year's publications can be considered as a response to the "necessity calls" [250,245,251,252].

Our review aims to contribute toward attention to the unifying direction for the studies of development, function, and decline of human perception, cognitive function, and action [252,253].

Article Information

Conflict of Interest Disclosures: None

Funding: None

Keywords

Stereochemistry; molecular biology, nonequilibrium biochemistry, phase transition, chirality transfer, chiral catalysis, protein folding, protein aggregation, neuronal cell, information processing, cognitive laterality; cognitive association.

References

- Dzygiel P & Wieczorek P. Stereo-selective Transport of Amino Acids and Peptides through Liquid Membranes. *Chemical Papers* 2002; 56:24-31.
- Alberts B, Johnson A, Lewis J, et al. *Molecular Biology of the Cell*. 4th edition. New York: Garland Science; 2002.
- St-Jean M and Sygusch J. Stereospecific Proton Transfer by a Mobile Catalyst in Mammalian Fructose-1, 6-bisphosphate Aldolase. *The Journal of Biological Chemistry* 2007;282:31028-31037.
- Solntsev KM, Bartolo EA, Pan G, et al. Excited-State Proton Transfer in Chiral Environments: Photo-racemization of BINOLs1. *Israel journal of chemistry* 2009;49:27-233.
- Patterson D and Schnell M. New studies on molecular chirality in the gas phase: enantiomer differentiation and determination of enantiomeric excess. *Physical Chemistry Chemical Physics* 2014;23.
- Fujiki M. Review. Supramolecular Chirality: Solvent Chirality Transfer in Molecular Chemistry and Polymer Chemistry. *Symmetry* 2014;6:677-703.
- Tiwari MP and Prasad A. Molecularly imprinted polymer based enantioselective sensing devices: a review. *Analytica chimica acta* 2015;853:1-18.
- Inaki M, Liu J and Matsuno K. Cell chirality: its origin and roles in left-right asymmetric development. *Philosophical Transactions of the Royal Society B* 2016;371:1710.
- Deng L, Wu S, Yao M and Gao C. Surface-anchored poly (acryloyl-L (D)-valine) with enhanced chirality-selective effect on cellular uptake of gold nanoparticles. *Scientific Reports* 2016;6:31595.
- McDowell G, Rajadurai S and Levin M. From cytoskeletal dynamics to organ asymmetry: a nonlinear, regulative pathway underlies left-right patterning. *Philosophical Transactions B* 2016;371:1710.
- Mousanezhada D, Haghpanaha B, Ghosha R, et al. Elastic properties of chiral, anti-chiral, and hierarchical honeycombs: A simple energy-based approach. *Theoretical and Applied Mechanics Letters* 2016;6:81-96.
- Morrow SM, Bisette AJ and Fletcher SP. Transmission of chirality through space and across length scales. *Nature Nanotechnology* 2017;12:410-419.
- Noyori R. Nobel lecture. Asymmetric Catalysis; science and opportunities 2001.
- Le Guennec P. On the concept of chirality. *Journal of Mathematical Chemistry* 1998;23:429-439.
- Ni J, Wang C, Zhang C, Hu Y, et al. Three-dimensional chiral microstructures fabricated by structured optical vortices in isotropic material. *Cornell Univ. Library* (205). ArXiv: 1608.01220 [physics.optics].
- Book by Nandi N. *Chirality in Biological Nanospaces: Reactions in Active Sites*. CRC Press. Taylor & Francis Group 2012.
- Piconese S, Tronelli G, Pippia P and Migliaccio F. Chiral and non-chiral nutations in Arabidopsis roots growth on the random positioning machine. *Journal of Experimental Botany* 2003;54:1909-1918.
- Cai H, Manoussaki D and Chadwick R. Effects of coiling on the micro-mechanics of the mammalian cochlea. *J R Soc Interface* 2005; 2: 341-348.
- Shannahoff-Khalsa D. Review. Lateralized rhythm of the central and autonomic nervous systems. *International Journal of Psychopathology* 1991;11:225-251.
- Novoselova NY and Saprionov NS. Role of inversion of interhemispheric asymmetry of phospholipid content in rat brain synaptosomes under stress conditions. *Doklady Biological Sciences* 2012; 442:7-10.
- Novoselova NY, Saprionov NS, Reichardt BA. Phospholipids of synaptosomes from rat brain hemispheres in dynamics of experimental myocardial infarction. Key role of right hemisphere in pathogenesis of depression associated with myocardial infarction (hypothesis). *Journal of Asymmetry* 2014;8:34-43.
- Weatherly CA, Du S, Parpia C, Santos PT, Hartman AL and Armstrong DW. d-Amino Acid Levels in Perfused Mouse Brain Tissue and Blood: A Comparative Study. *ACS Chemical Neuroscience* 2017;8:1251-1261.
- GM, Mao B and Cho KC. Chiral Features of Proteins. In the book by Paul G. Mezey: *New Developments in Molecular Chirality* 1991.
- Tan HH, Makino A, Sudesh K, Greimel P and Kobayashi T. Spectroscopic evidence for the unusual stereochemical configuration of an endosome-specific lipid. *Angewandte Chemie International Edition* 2012;51:533-535.
- Book by Würges K. *Enzyme Supported Crystallization of Chiral Amino Acids*. Forschungszentrum Jülich 2011.
- Satir P. Chirality of the cytoskeleton in the origins of cellular asymmetry. *Philosophical Transactions of the Royal Society B* 2016;371.
- Li M, Zhao C, Ren J and Qu X. Chiral Metallo-Supramolecular Complex Directed Enantioselective Self-Assembly of β -Sheet Breaker Peptide for Amyloid Inhibition. *Small* 2015;11:4651-4655.
- Scala CD, Yahi N, Boutemour S, et al. Common molecular mechanism

- of amyloid pore formation by Alzheimer's β -amyloid peptide and alpha-synuclein. *Scientific Reports* 2016;6:28781.
29. Thakur GA, Palmer SL, Harrington PE, et al. Enantiomeric resolution of a novel chiral cannabinoid receptor ligand. *Journal of Biochemical and Biophysical Methods* 2002; 54:415-422.
 30. Book edited by Eichelbaum MF, Testa B and Somogyi A. *Stereochemical Aspects of Drug Action and Disposition*. Springer-Verlag Berlin Heidelberg (2003).
 31. Nguyen LA, He H and Pham-Huy C. Chiral Drugs: An Overview. *International Journal of Biomedical Science* 2006;2:85-100.
 32. Hutt AJ and O'Grady J. Drug chirality: a consideration of the significance of the stereochemistry of antimicrobial agents. *Journal of Antimicrobial Chemotherapy* 1996;37:7-32.
 33. Dixit S and Hag Park J. Application of antibiotics as chiral selectors for capillary electrophoretic enantioseparation of pharmaceuticals: a review. *Biomedical Chromatography* 2014;28:10-26.
 34. Wächtershäuser G. From pre-cells to Eukarya – a tale of two lipids. *Molecular Microbiology* 2003;47:13-22.
 35. Weiskopf RB, Nau C and Strichartz GR. Drug Chirality in Anesthesia. *Anesthesiology* 2002;97:497-502.
 36. Nanda V, Andrianarijaona A and Narayanan C. The role of protein homochirality in shaping the energy landscape of folding. *Protein Science: a publication of the Protein Society* 2007;8:1667-1675.
 37. Ali HS, Alhaj OA, Al-Khalifa AS and Brückner H. Determination and stereochemistry of proteinogenic and non-proteinogenic amino acids in Saudi Arabian date fruits. *Amino Acids* 2014;46:2241-2257.
 38. Drechsel-Grau C and Marx D. Tunnelling in chiral water clusters: Protons in concert. *Nature Physics* 2015;11:216-218.
 39. Matsumoto Y, Chen R, Anikeeva P and Jasanoff A. Engineering intracellular biomineralization and biosensing by a magnetic protein. *Nature Communications* 2015;6:8721.
 40. Spector MS, Easwaran KRK, Jyothi G, et al. Chiral molecular self-assembly of phospholipid tubules: A circular dichroism study. *Proceedings of the National Academy of Sciences of the United States of America* 1996;93:12943-12946.
 41. S. Bisht S, Pandey P, Bhargava B, et al. Bioremediation polyaromatic hydrocarbons (PAHs) using rhizosphere technology. *Brazilian Journal of Microbiology* 2015;46:7-21.
 42. Kessler D, Kallenbach M, Diezel C, et al. How scent and nectar influence floral antagonists and mutualists. *eLife* 2015;4:e07641.
 43. Petersen P. I. Speaker. Water at surfaces with tunable surface chemistries and the chiral imprint of water around DNA. *APS March Meeting 2016*. Vol. 61, Number 2. Monday-Friday, March 14-18, 2016; Baltimore, Maryland. Session P31: Water at Interfaces: From Spectroscopy Techniques to Computer Simulations. P31.00001 (March 16, 2016).
 44. Hazen RM. Mineral surfaces, geochemical complexities, and the origin of life. *Cold Spring Harbor perspectives in biology* 2010;2:a002162.
 45. Blackmond DG. The origin of biological homochirality. *Cold Spring Harbor perspectives in biology* Labor. Press. 2010.
 46. Malyshko EV and Tverdislov VA. Chirality as a physical aspect of structure formation in biological macromolecular systems. *Journal of Physics: Conference Series* 2016;741:1.
 47. Chhabra SR, Harty C, Hooi DS, Daykin M, et al. Synthetic analogues of the bacterial signal (quorum sensing) molecule N-(3-oxododecanoyl)-L-homoserine lactone as immune modulators. *Journal of medicinal chemistry* 2003;46:97-104.
 48. Yan ECY, Fu L, Wang Z and Liu W. Review. Biological Macromolecules at Interfaces Probed by Chiral Vibrational Sum Frequency Generation Spectroscopy. *Chemical Reviews* 2014;114:8471-8498.
 49. Book edited by Lichtfouse E, Schwarzbauer J and Robert D. *Environmental Chemistry for a Sustainable World: Volume 2: Remediation of Air and Water Pollutions*. Springer Sc. +Bus. Media B.V. 2012.
 50. Martin M and Jones GV. Motor imagery theory of contralateral handedness effect in recognition memory: toward a chiral psychology of cognition. *Journal of Experimental Psychology, General*. 1999;128:265-282.
 51. Burke WJ and Kratochvil ChJ. Stereoisomers in psychiatry: the case of escitalopram. *Journal of clinical psychiatry* 2002;4:20-24.
 52. Baker GB, Prior TI, and Coutts RT. Chirality and drugs used to treat psychiatric disorders. *Journal of psychiatry and neuroscience* 2002;6:401-403.
 53. Patocka J and Dvorak A. Biomedical aspects of chiral molecules. *Journal of Applied Biomedicine* 2004;2:95-100.
 54. Zawirska-Wojtasiak R. Chirality and the Nature of Food Authenticity of Aroma. *Acta Scientiarum Polonorum, Technologia Alimentaria* 2006;5: 21-36.
 55. Carlavilla D, Victoria Moreno-Arribas M, Fanali S, Cifuentes A. Chiral MEKC-LIF of amino acids in foods: Analysis of vinegars. *Electrophoresis* 2006;27:2551-2557.
 56. Li B and Haynie DT. Chiral Drug Separation. In the book edited by Sunggyu Lee. *Encyclopedia of Chemical Processing* 2006;1.
 57. Wang P, Yu SJ, Govorov AO and Ouyang M. Cooperative expression of atomic chirality in inorganic nanostructures. *Nature Communications* 2017.
 58. Ariga K, Richards GJ, Ishihara S, Izawa H and Hill JP. Review. Intelligent Chiral Sensing Based on Supramolecular and Interfacial Concepts. *Sensors* 2010;10:6796-6820.
 59. Prelog V. Chirality in Chemistry. Nobel Lecture, December 12, 1975. The Nobel Foundation (1975). From Nobel Lectures, Chemistry 1971-1980, Editor-in-Charge Tore Frängsmyr, Editor Sture Forsén, World Scientific Publishing Co., Singapore, 1993.
 60. Frank. On spontaneous asymmetric synthesis. *Biochimica et biophysica acta* 1953;4:459-63.
 61. Jafarpour F, Biancalani T, Goldenfeld N. Noise-induced symmetry breaking far from equilibrium and the emergence of biological homochirality. *Physical Review E* 2017;95(3-1):032407.
 62. Ulbricht TLV and Vester F. Attempts to induce optical activity with polarized β -radiation. *Tetrahedron* 1962;18:629-637.
 63. Dreiling JM and Gay TJ. Chirality Sensitive Electron-Induced Molecular breaking and the Vester-Ulbricht Hypothesis. *Physical Review Letters* 2014;113:118103.
 64. Dreiling and Joan. Chiral Sensitivity in Electron-Molecular Interactions. *APS Gaseous Electronics Conference 2015*, abstract #IW4.001The Smithsonian/NASA Astrophysics Data System.
 65. Viedma C. Chiral symmetry breaking crystallization: complete chiral purity induced by nonlinear autocatalysis recycling. *Physical Review Letters* 2005;94:065504.
 66. Klussmann M, Iwamura H, Mathew SP, Wells DH, et al. Thermodynamic control of asymmetric amplification in amino acid catalysis. *Nature* 2006;441:621-623.
 67. Blackmond DG. Chiral amnesia as a driving force for solid-phase homochirality. *Chemistry European. Journal*. 2007;13:3290-3295.
 68. Viedma C. Chiral symmetry breaking and complete chiral purity in thermodynamic-kinetic feedback near equilibrium: implication for the origin of biochirality. *Astrobiology* 2007;7:312-319.
 69. Cronin J and Reisse J. Chirality and the Origin of Homochirality. Ch. 3 in the Book edited by Muriel Gargaud, Bernard Barbier, Hervé Martin, Jacques Reisse. *Lectures in Astrobiology*. 1:473. Edited by Muriel Gargaud, Bernard Barbier, Hervé Martin and Jacques Reisse.
 70. Blackmond DG. The origin of biological homochirality. *Cold Spring Harbor perspectives in biology* 2010;2:a002147.
 71. McGuire BA, Carroll PB, Loomis RA, et al. Discovery of the Interstellar Chiral Molecule Propylene Oxide (CH₃CHCH₂O). *Science* 2016;352:1449-1452.
 72. Hein JE and Blackmond DG. On the origin of single chirality of amino acids and sugars in biogenesis. *Accounts of chemical research*.

- 2012;45:2045–2054.
73. Yin P, Zhang ZM, Lv H, et al. Chiral Recognition and Selection during the Self-Assembly Process of Protein-Mimic Macroanions. *Nature Communications* 2015;6:6475.
 74. Bouvier M. Review. Oligomerization of G-protein-coupled transmitter receptors. *Nature Reviews Neuroscience* 2012;2:274-286.
 75. Turner NJ. Controlling chirality. *Current Opinion in Biotechnology* 2003;14:401-416.
 76. Kanga L and Lubenskya TC. Chiral twist drives raft formation and organization in membranes composed of rod-like particles. *Proceedings of the National Academy of Sciences of the United States of America* 2017;114:E19–E27.
 77. Pemberton TA and Christianson DW. Review. Genera base-general acid catalysis by terpenoid cyclases. *The Journal of Antibiotics* 2016;69:486–493.
 78. Shiho K, Kumatabara Y and Shirakawa S. A new generation of chiral phase-transfer catalysts. *Organic and Biomolecular Chemistry* 2016;14:5367-5376.
 79. Stamatopoulos P, Brohan E, Prevost C, et al. Influence of Chirality of Lactones on the Perception of Some Typical Fruity Notes through Perceptual Interaction Phenomena in Bordeaux Dessert Wines. *Journal of agricultural and food chemistry* 2016;64:8160-8167.
 80. Book edited by Stamenov M and Gallese V. *Mirror Neurons and the Evolution of Brain and Language*. John Benjamins B.V. 2002.
 81. Pizzarello S. The chemistry of life's origin: A carbonaceous meteorite perspective. *Accounts of Chemical Research* 2006;39:231-237.
 82. Zahnle K, Schaefer L and Fegley B. Earth's early atmospheres. *Cold Spring Harbor perspectives in biology* 2010;2:a004895.
 83. Meinert C. Understanding the Origin of Chiral Molecules in Interstellar Space. *Séminaires LLR. Institut de chimie de Nice. (Europe/Paris)* 2016.
 84. Singh D. Neutrino helicity and chirality transitions in Schwarzschild space-time. *Physical Review D* 2005;71.
 85. Sarasij RC and Rao M. Tilt texture domains on a membrane and chirality induced budding. *Physical Review Letters* 2002;88.
 86. Lin SK. The Nature of the Chemical Process. 1. Symmetry Evolution - Revised Information Theory, Similarity Principle and Ugly Symmetry. *International Journal of Molecular Sciences* 2001;2:10-39.
 87. Wang Y and Nathans J. REVIEWS. Tissue/planar cell polarity in vertebrates: new insights and new questions. *Development* 2007;134:647-658.
 88. Lawrence PA, Struhl G and Casal J. Perspective Opinion. Planar cell polarity: one or two pathways? *Nature Reviews Genetics* 2007;8:555-563.
 89. Vandenberg LN and Levin M. A unified model for left-right asymmetry? Comparison and synthesis of molecular models of embryonic laterality. *Developmental biology* 2013;379:1-15.
 90. Axelrod JD. Strabismus comes into focus. *Nature Cell Biology* 2002;4:E6-E8.
 91. Le Bel JA. "Sur les relations qui existent entre les formules atomiques des corps organiques et le pouvoir rotatoire de leurs dissolutions" (On the relations that exist between the atomic formulas of organic substances and the rotatory power of their solutions), *Bulletin de la Société Chimique de Paris* 1874;22:337–347.
 92. van't Hoff JH. A suggestion looking to the extension into space of the structural formulas at present used in chemistry. And a note upon the relation between the optical activity and the chemical constitution of organic compounds. *Archives neerlandaises des sciences exactes et naturelles* 1874; 9: 445-454.
 93. Weisberg M, Needham P and Hendry R. *Philosophy of Chemistry*. Stanford Encyclopedia of Philosophy 2011.
 94. Wolf C. *Dynamic Stereochemistry of Chiral Compounds: Principles and Applications*. Copyright: 2007.
 95. Tsukube H and Shinoda S. Lanthanide complexes in molecular recognition and chirality sensing of biological substrates. *Chemical Reviews* 2002;102:2389–2404.
 96. Song G and Ren J. Recognition and regulation of unique nucleic acid structures by small molecules. *Chemical Communications* 2010;46:7283–7294.
 97. J. Crassous J. Transfer of chirality from ligands to metal centers: Recent examples. *Chemical Communications* 2012;48:9684-9692.
 98. Kolomiets E, Berl V and Lehn JM. Chirality induction and protonation-induced molecular motions in helical molecular strands. *Chemistry A European Journal* 2007;13:5466-5479.
 99. Eerdun C, Hisanaga S and Setsune J. Single helicates of dipalladium (II) hexapyrroles: Helicity induction and redox tuning of chiroptical properties. *Angewandte Chemie International Education* 2013;52:929-932.
 100. Brown RA, Diemer V, Webb SJ, Clayden J. End-to-end conformational communication through a synthetic purinergic receptor by ligand-induced helicity switching. *Nature Chemistry* 2013;5:853-860.
 101. Miyake H. *Supramolecular Chirality in Dynamic Coordination Chemistry*. *Symmetry* 2014;6:880-895.
 102. K. Soai K. *Amplification of Chirality*. Pg. 73. *Topic in Current Chemistry* 2008;284. Springer-Verlag Berlin (2008).
 103. Ito H, Ikeda M, Hasegawa T, Furusho Y and Yashima E. Synthesis of complementary double-stranded helical oligomers through chiral and achiral amidinium-carboxylate salt bridges and chiral amplification in their double-helix formation. *Journal of the American Chemical Society* 2011;133:3419-3432.
 104. Yashima E, Ousaka N, Taura D, et al. *Supramolecular Helical Systems: Helical Assemblies of Small Molecules, Foldamers, and Polymers with Chiral Amplification and Their Functions*. *Chemical Reviews* 2016;116:13752-13990.
 105. Kafri R, Markovitch O and Lancet D. Spontaneous chiral symmetry breaking in early molecular networks. *Biology Direct* 2010;5:38.
 106. Li Y, Wang M, Li Q, et al. Azoarenes with opposite chiral configuration: light-dravercible handedness inversion in self-organized helical superstructures. *Angewandte Chemie International Edition* 2013;52:8925-8929.
 107. Li Q. *Nanoscience with Liquid Crystals: From Self-Organized Nanostructures to Applications*. Springer Int. Publish. Switzerland 2014.
 108. Jiang X, Lim YK, Zhang BJ, et al. Dendritic molecular switch: Chiral folding and helicity inversion. *Journal of the American Chemical Society* 2008;130:16812-16822.
 109. Akine S, Hotate S and Nabeshima T. A molecular leverage for helicity control and helix inversion. *Journal of the American Chemical Society* 2011;133:13868-13871.
 110. Miyake H, Ueda M, Murota S, Sugimoto H and Tsukube H. Helicity inversion from left- to right-handed square planar Pd (II) complexes: Synthesis of a diastereomer pair from a single chiral ligand and their structure dynamism. *Chemical Communications* 2012;48:3721-3723.
 111. Weigelt S, Busse C, Petersen L, et al. Chiral switching by spontaneous conformational change in adsorbed organic molecules. *Nature Materials* 2006;5:112-117.
 112. Pijper D, Jongejan MGM, Meetsma A and Feringa BL. Light-controlled supramolecular helicity of a liquid-crystalline phase using a helical polymer functionalized with single chiroptical molecular switch. *Journal of the American Chemical Society* 2008;130:4541-4552.
 113. Stillinger FC. *Microscopic Kinetic Model Exhibiting Chiral Symmetry Breaking*. Bruce Berne Symposium, ACS, Boston, Aug. 22-25, 2010.
 114. Hajlaouia F, Yahyaouia S, Nailia H, Mhiria T and Bataille T. Crystal structure, phase transition and thermal decomposition of a copper (II) sulfate dihydrate containing a chiral organic ammonium cation. *Inorganica Chimica Acta* 2010;363:691-695.
 115. Dierking I. *Chiral Liquid Crystals: Structure, Phases, Effects*. *Symmetry* 2014;6:444-472.

116. Smith E and Morowitz HJ. *The Origin and Nature of Life on Earth: The Emergence of the Fourth Geosphere*. Cambridge University Press 2016.
117. Butler MT and John B. Wallingford. Planar cell polarity in development and disease. *Nature Reviews Molecular Cell Biology* 2017;18:375-388.
118. Strutt D and Goodrich LV. Review. Principles of planar polarity in animal development. *Development* 2011;138:1877-1892.
119. Chela-Flores J. The origin of chirality in protein amino acids. *Chirality* 1994;6:165-168.
120. Iachello F. Analytic Description of Critical Point Nuclei in a Spherical-Axially Deformed Shape Phase Transition. *Physical Review Letters* 2001;87.
121. Alloeyau D, Ricolleau C, Mottet C, et al. Size and shape effects on the order-disorder phase transition in CoPt nanoparticles. *Nature Materials* 2009;8:940-946.
122. Nishiyama I, Yamamoto J, Goodby JW and Yokoyama H. Chiral smectics: Molecular design and superstructures. *Molecular Crystals and Liquid Crystals* 2005;443:25-41.
123. Callister T, Sammut L, Qiu S, Mandel I and Thrane E. Limits of Astrophysics with Gravitational-Wave Backgrounds. *Physical Review X* 2016;6.
124. Wang JS, Feng XQ, Xu J, Qin QH and Yu SW. Chirality Transfer from Molecular to Morphological Scales in Quasi-One-Dimensional Nanomaterials: A Continuum Model. *Journal of Computational and Theoretical Nanoscience* 2011;8:1278-1287.
125. Qing G and Sun T. The transformation of chiral signals into macroscopic properties of materials using chirality-responsive polymers. *NPG Asia Materials* 2012;4:e4.
126. De Michele C, Zanchetta G, Bellini T, Frezza E and Ferrarini A. Hierarchical Propagation of Chirality through Reversible Polymerization: The Cholesteric Phase of DNA Oligomers. *ACS Macro Letters* 2016;5:208-212.
127. Couillet P, Lega J, Houchmandzadeh B and Lajzerowicz J. Breaking chirality in nonequilibrium systems. *Physical Review Letters* 1990;65:1352.
128. Tjhunga E, Catesa ME and Marenduzzob Dd. Contractile and chiral activities codetermine the helicity of swimming droplet trajectories. *Proceedings of the National Academy of Sciences of the United States of America* 2017;118:4631-4636.
129. Pi-Boleda B, Sans M, Campos M, et al. Studies on Cycloalkane-Based Bisamide Organogelators: A New Example of Stochastic Chiral Symmetry-Breaking Induced by Sonication. *A European Journal of Chemistry* 2017;23:3357-3365.
130. Wang JS, Wang G, Feng XQ, et al. Hierarchical chirality transfer in the growth Towel Gourd tendrils. *Scientific Reports* 2013;3:3102.
131. Pasteur L. Sur les corpuscles organizes qui existent dans l'atmosphère: Examen de la doctrine des generations spontanées. *Leçon Professe à la Société Chimique de Paris*. 19 Mai 1861.
132. Gal J. Louis Pasteur, Language, and Molecular Chirality. I. Background and Dissymmetry. *Chirality* 2011;23:1-16.
133. Mason SF. *Molecular Optical Activity and the Chiral Discriminations*. Cambridge University Press. 1982.
134. Bonet JC, Ignés-Mullol J and Sagués F. Chiral selection under swirling: From a concept to its realization in soft-matter self-assembly. *Contributions to science* 2015;11:199-205. Thematic issue on Non-equilibrium physics.
135. Kozlova SG, Kompankov NB and Zavakhina MS. Activation Parameters of Self-Diffusion of Aromatic Chiral Molecules in External Magnetic Fields. *Journal of Physical Chemistry B*. 2017;121:6655-6658.
136. Bonner WA. Chirality and life. *Origins of Life and Evolution of the Biosphere* 1995;25:175-190.
137. Shimada A, Ozaki H, Saito T and Noriko F. Tryptophanase-Catalyzed L-Tryptophan Synthesis from d-Serine in the Presence of Diammonium Hydrogen Phosphate. *International journal of molecular sciences* 2009;10:2578-2590.
138. Jeong J, Aetukuri N, Graf T, et al. Suppression of Metal-Insulator Transition in VO₂ by Electric-Field-Induced Oxygen Vacancy Formation. *Science* 2013; 339: 402-1405.
139. Palyi G, Zucchi C and Caglioti L. *Progress in Biological Chirality*. Chapter: 26. Shimada A, Fujii N and Saiko T. Tryptophanase on D-Tryptophan. Introduction. Elsevier Ltd (2004).
140. Barron LD. Chirality: spin and gravity give a helping hand. *Nature Chemistry* 2012;4:150-152.
141. Ousaka N, Takeyama Y, Iida H and Yashima E. Chiral information harvesting in dendritic metalloptides. *Nature Chemistry* 2011;3:856-861.
142. Albrecht M. Let's twist again—double-stranded, triple-stranded, and circular helicates. *Chemical Reviews* 2001;101:3457-3498.
143. Clayden J. Molecular devices: Communicating chirality. *Nature Chemistry* 2011;3:842-843.
144. Palmans ARA and Meijer EW. Amplification of chirality in dynamic supramolecular aggregates. *Angewandte Chemie International Edition* 2007;46:8948-8968.
145. Aw S, and Levin M. Hypothesis. Is left-right asymmetry a form of planar cell polarity? *Development* 2009;136:355-366.
146. Kannan S and Zacharias M. Folding simulations of Trp-cage mini protein in explicit solvent using biasing potential replica-exchange molecular dynamics simulations. *Proteins* 2009;76:448-460.
147. Jacobsen EN, Pfaltz A and Yamamoto H. *Comprehensive Asymmetric Catalysis*. Springer, Berlin. 1999; I-III.
148. Ojima I. *Catalytic Asymmetric Synthesis*. Wiley, New York, 3rd Edition 2000.
149. Hosamani B, Ribeiro MF, da Silva Júnior EN and Namboothiri NNI. Catalytic asymmetric reactions and synthesis of quinones. *Journal of Organic & Biomolecular Chemistry* 2016;29: 6913-6931.
150. Faraday M. Esq. D.C.L. F.R.S. *LIV. Thoughts on Ray-vibrations*. *Philosophical Magazine* 1846;28.
151. Dvornikov M and Semikoz V. Lepton asymmetry growth in the symmetric phase of an electroweak plasma with hypermagnetic fields versus its washing out by sphalerons. *Physical Review* 2013.
152. Woszczyk A and Szabelski P. Enhancing the separation of enantiomers in absorbed overlayers: a Monte Carlo study. *Annales Universitatis Mariae Curie-Skłodowska. Lublin - Polonia*. 2014;68:133-141.
153. Volovik GE and Krusius M. Viewpoint: Chiral Quantum Textures. *Physics* 2012;5:130.
154. Kessler EM, Giedke G, Imamoglu A, et al. Dissipative phase transition in a central spin system. *Physical Review A* 2012;86.
155. Krokhamalskii T, Baliha V, Derzhko O, Schulenburg J and Richter J. Frustrated honeycomb-lattice bilayer quantum antiferromagnet in a magnetic field: Unconventional phase transitions in a two-dimensional isotropic Heisenberg model. *Physical Review B* 2017;95.
156. Labuta J, et al. NMR Spectroscopic detection of chirality and enantiopurity in referenced systems without formation of diastereomers. *Nature Communications* 2013;4: 2188.
157. Pasteur L. C.r. hebd. Séanc. Acad. Sci. Paris. 1848. 26;535. *Oeuvres de Pasteur Vol. 1* (ed. Pasteur Valéry-Radot) 61-64 (Masson, Paris, 1922).
158. Parker D. NMR determination of enantiomeric purity. *Chemical Reviews* 1991;91:1441-1457.
159. Park J, Goh M and Akagi K. Helical Nylons and Polyphthalamides Synthesized by Chiral Interfacial Polymerizations between Chiral Nematic Liquid Crystal and Water Layers. *Macromolecules* 2014;47:2784-2795.
160. Lodah P, Mahmoodian S, Stobbe S, et al. Chiral quantum optics. *Nature* 2017;541:473-480.

161. Ariga K, Vinu A, Hill JP, Mori T. Coordination chemistry and supramolecular chemistry in mesoporous nanospace. *Coordination Chemistry Reviews* 2007;251:2562-2591.
162. Nakanishi T, Michinobu T, Yoshida K, et al. Nanocarbon superhydrophobic surfaces created from fullerene-based hierarchical supramolecular assemblies. *Advanced Materials* 2008;20:443-446.
163. Nakanishi T, Takahashi H, Michinobu T, et al. Fine-tuning supramolecular assemblies of fullerenes bearing long alkyl chains. *Thin Solid Films* 2008;516:2401-2406.
164. Okamoto K, Chithra P, Richards JG, JP, Ariga K. Self-assembly of optical molecules with supramolecular concepts. *International journal of molecular sciences* 2009;10:1950-1966.
165. Mandal S, Lee MV, Hill JP, Vinu A and Ariga K. Recent developments in supramolecular approach for nanocomposites. *Journal of Nanoscience and Nanotechnology* 2010;10:21-33.
166. Pu L. Fluorescence of organic molecules in chiral recognition. *Chemical Reviews* 2004;104:1687-1716.
167. Watarai H and Adachi K. Measuring the optical chirality of molecular aggregates at liquid-liquid interfaces. *Analytical and Bioanalytical Chemistry* 2009;395:1033-1046.
168. Xiao W, Ernst KH, Palotas K, et al. Microscopic origin of chiral shape induction in achiral crystals. *Nature Chemistry* 2016; 8:326-330.
169. Fu W and Tang W. Chiral Monophosphorus Ligands for Asymmetric Catalytic Reactions. *ACS Catalysis* 2016;6: 4814-4858.
170. Govana J and Gun'ko YK. Recent progress in chiral inorganic nanostructures. *Nanoscience* 2016;3:1-30.
171. Ben-Moshe A, Maoz BM, Govorov AO and Markovich G. Chirality and chiroptical effects in inorganic nanocrystal systems with plasmon and exciton resonances. *Chemical Society Reviews*. 2013;42:7028-7041.
172. Pagni RM. The weak nuclear force, the chirality of atoms, and the origin of optically active molecules. *Foundations of Chemistry* 2009;11:105-122.
173. Kawagoe Y, Fujiki M and Nakanoa Y. Limonene magic: noncovalent molecular chirality transfer leading to ambidextrous circularly polarised luminescent π -conjugated polymers. *New Journal of Chemistry* 2010;34:637-647.
174. Brown NA and Wolpert L. The development of handedness in left/right asymmetry. *Development* 1990;109:1-9.
175. Capozziello S and Lattanzi A. Spiral galaxies as enantiomers: Chirality, an underlying feature in chemistry and astrophysics. *Chirality* 2006;18:17-23.
176. Boyd R. *Chiral Stardust, Supernovae and the Molecules of Life: Might We All Be Aliens?* Springer Science+Business Media. 2012.
177. Machleidt R and Entem DR. Chiral effective field theory and nuclear forces. *Physics Reports* 2011;503:1-75.
178. Vasseur DA and Fox JW. Phase-locking and environmental fluctuations generate synchrony in a predator-prey community. *Letter Nature* 2009;460:1007-1010.
179. Lin H, Jiang Y, Zhang Q, Hua K and Li Z. An in-tether sulfilimine chiral center induces helicity in short peptides. *Chemical Communications* 2016;52:10389-10391.
180. Letokhov VS. On difference of energy levels of left and right molecules due to weak interactions. *Physics Letters A* 1975;53:275-276.
181. Quick M. How important is parity violation for molecular and bimolecular chirality? *Angewandte Chemie International Edition* 2002;41:4618-4630.
182. Xu K. Anisotropic 2s2p Orbitals as Simple Descriptors of the Chirality of Carbon Centres. *Australian Journal of Chemistry* 2016;69:775-784.
183. Jackson G. Types of Biological Macromolecules.
184. Kwiecińska J and Cieplak M. Chirality and protein folding. *Journal of Physics: Condensed Matter* 2005;17.
185. Stefani M. Protein Folding and Misfolding on Surfaces. *International Journal of Molecular Sciences* 2009;9:2515-2542.
186. Onuchic JN, Luthey-Schulten Z and Wolynes PG. Theory of protein folding: the energy landscape perspective. *Annual Review of Physical Chemistry* 1997;48:545-600.
187. Duan Y and Kollman PA. Pathways to a protein folding intermediate observed in a 1-microsecond simulation in aqueous solution. *Science* 1998;282:740-744.
188. Daggett V and Fersht A. The present view of the mechanism of protein folding. *Nature Reviews Molecular Cell Biology* 2003;4:497-502.
189. Kubelka J, Hofrichter J and Eaton WA. The protein folding 'speed limit'. *Current Opinion in Structural Biology* 2004;14:76-88.
190. Seibert MM, Patriksson A, Hess B and van der Spoel D. Reproducible polypeptide folding and structure prediction using molecular dynamics simulations. *Journal of Molecular Biology* 2005;354:173-183.
191. Scheraga HA, Khalili M and Liwo A. Protein-folding dynamics: Overview of molecular simulation techniques. *Annual Review of Physical Chemistry* 2007;58:57-83.
192. Dill KA, Ozkan SB, Weikl TR, Chodera JD and Voelz VA. The protein folding problem: When will it be solved? *Current Opinion in Structural Biology* 2007;17:342-346.
193. Ensign DL, Kasson PM and Pande VS. Heterogeneity even at the speed limit of folding: Large-scale molecular dynamics study of a fast-folding variant of the villin headpiece. *Journal of Molecular Biology* 2007;374:806-816.
194. Juraszek J, Bolhuis PG. Rate constant and reaction coordinate of Trp-cage folding in explicit water. *Biophysical Journal* 2008;95:4246-4257.
195. Freddolino PL and Schulten K. Common structural transitions in explicit-solvent simulations of villin headpiece folding. *Biophysical Journal* 2009;97:2338-2347.
196. Zhanga C and Maa J. Folding helical proteins in explicit solvent using dihedral-biased tempering. *Proceedings of the National Academy of Sciences of the United States of America* 2012;109:8139-8144.
197. Sarasij RC, Mayor S and Rao M. Chirality-Induced Budding: A Raft-Mediated Mechanism for Endocytosis and Morphology of Caveolae? *BioPhysical Journal* 2007;92:3140-3158.
198. Ho RM, Wang HF and Li MC. Chirality Transfer in Chiral Polymers and Block Copolymers. *Encyclopedia of Polymer Science and Technology* 2014.
199. Cabeen MT and Jacobs-Wagner C. Bacterial cell shape. *Nature Reviews Microbiology* 2005;3:601-610.
200. Gitai Z. The new bacterial cell biology: Moving parts and subcellular architecture. *Cell* 2005;120:577-586.
201. Gaballah A, Kloeckner A, Otten C, Sahl HG and Henrichfreise B. Functional Analysis of the Cytoskeleton Protein MreB from *Chlamydomonas pneumoniae*. *PLoS ONE* 2011;6:e25129.
202. Wang S, Furchtgott L, Huang KC and Shaevitz JW. Helical insertion of peptidoglycan produces chiral ordering of the bacterial cell wall. *Proceedings of the National Academy of Sciences of the United States of America* 2012;109(10): E595-E604.
203. Hatori R, Ando T, Sasamura T, et al. Left-right asymmetry is formed in individual cells by intrinsic cell chirality. *Mechanisms of Development* 2014;133:146-62.
204. Zhong-can OY and Jixing L. Theory of helical structures of tilted chiral lipid bilayers. *Physical Review A* 1991;43:6826-6836.
205. Eghiaian F. Lipid Chirality Revisited: A Change in Lipid Configuration Transforms Membrane-Bound Protein Domains. *Biophysical Journal* 2015;108:2757-2758.
206. Lister FGA, Le Bailly BAF, Webb SJ and Jonathan C. Ligand-modulated conformational switching in a fully synthetic membrane-bound receptor. *Nature Chemistry* 2017;9:420-425.
207. Darwin C. On the movement and habits of climbing plants. *John Murray. London. 1865.*
208. Isnard S and Silk WK. Moving with climbing plants from Charles Darwin's time into 21st century. *American Journal Botany* 2009;96:1205-1221.

209. Ye HM, et al. Surface stress effects on the bending direction and twisting chirality of lamellar crystals of chiral polymer. *Macromolecules* 2010;43:5762-5770.
210. Grande C & Patel NH. Nodal signaling is involved in left-right asymmetry in snails. *Nature* 2009;475:1007-1011.
211. Lough WJ and Wainer IW. Chirality in mature and applied science. Blacjwel Science, Oxford 2002.
212. Qiu D, Cheng SM, Wozniak L, et al. Localization and loss-of-function implicate ciliary proteins in early, cytoplasmic LR asymmetry. *Developmental Dynamics* 2005;234:176-189.
213. Pediconi MF, de Fernández AMR and Barrantes FJ. Asymmetric distribution and down-regulation of the muscarinic acetylcholine receptor in rat cerebral cortex. *Neurochemical research* 1993;18:565-72.
214. Pavlov IP. Conditional Reflexes. 1927/1960.
215. Galaburda AM, LeMay M, Kemper TL and Geschwind N. Right-left asymmetries in the brain. *Science* 1978;199:852-856.
216. Toga AW and Thompson PM. Mapping brain asymmetry. *Nature Reviews Neuroscience* 2003;4:37-48.
217. Kertesz A, Polk M, Black SE and Howell J. Sex, handedness, and the morphometry of cerebral asymmetries on magnetic resonance imaging. *Brain research* 1990;530:40-48.
218. Good CD, Johnsrude I, Ashburner J, et al. Cerebral asymmetry and the effects of sex and handedness on brain structure: a voxel-based morphometric analysis of 465 normal adult human brains. *Neuroimage* 2001;14:685-700.
219. Watkins KE, Paus T, Lerch JP, et al. Structural asymmetries in the human brain: a voxel-based statistical analysis of 142 MRI scans. *Cerebral Cortex* 2001;11:868-877.
220. Shaw P, Lalonde F, Lepage C, et al. Development of cortical asymmetry in typically developing children and its disruption in attention-deficit/hyperactivity disorder. *Archives of General Psychiatry* 2009;66:888-896.
221. Zhou D, Lebel C, Evans A and Beaulieu C. Cortical thickness asymmetry from childhood to older adulthood. *Neuroimage* 2013;83:66-74.
222. Lin A, Clasen L, Lee NR, et al. Mapping the Stability of Human Brain Asymmetry across Five Sex-Chromosome Aneuploidies. *Journal of Neuroscience* 2015;35:140-145.
223. Duyn JH. EEG-fMRI Methods for the Study of Brain Networks during Sleep. *Frontiers in Neurology* 2012;3:100.
224. Hughahl K and Davidson RJ. *The Asymmetrical Brain*. The MIT Press. 2003.
225. Hugdahl K and Westerhausen R. *The Two Halves of the Brain*. MIT Press. 2010.
226. Bryden M. *Laterality Functional Asymmetry in the Intact Brain*.
227. Mori K. Significance of chirality in pheromone science. *Bioorganic & Medicinal Chemistry* 2007;15:7505-7523.
228. Wyatt TD. *Pheromones and Animal Behavior: Chemical Signals and Signatures*. 2nd Edition Cambr. Univ. Press 2014.
229. de Moraes Júnior R, de Sousa BM and Fukusima S. Hemispheric specialization in face recognition: from spatial frequencies to holistic/analytic cognitive processing. *Psychology and Neuroscience* 2014;7:503-511.
230. Huber BA. Mating positions and the evolution of asymmetric insect genitalia. *Genetica* 2010;138:19-25.
231. Bakalkin GYa. Review Article. Neuropeptides induce directional asymmetry in brain and spinal cord: facts and hypotheses. *The International journal of neuroscience* 1989;48:105-124.
232. Dubovy SR, Fernandez MP, Echegaray JJ, et al. Expression of hypothalamic neurohormones and their receptors in the human eye. *Oncotarget* 2017. DOI: 10.18632/oncotarget.18358.
233. Wang J and Feringa BL. Dynamic control of chiral space in a catalytic asymmetric reaction using a molecular motor. *Science* 2011;331:1429-1432.
234. Kistemaker JCM, Štacko P, Visser J and Feringa BL. Unidirectional rotary motion in achiral molecular motors. *Nature Chemistry* 2015;7:890-896.
235. Silvi S, Ventura M and Credi A. Artificial molecular shuttles: from concepts to devices. *Journal of Materials Chemistry* 2009;19:2279-2294.
236. Zhu K, O'Keefe CA, Vukotic VN, Schurko RW and Loeb SJ. A molecular shuttle that operates inside a metal-organic framework. *Nature Chemistry* 2015;7:514-519.
237. Yamaki M, Nakayama S, Hoki K, Kono H and Fujimura Y. Quantum dynamics of light-driven chiral molecular motors. *Physical Chemistry Chemical Physics* 2009;11:1662-1678.
238. von Delius M and Leigh DA. Walking molecules. *Chemical Society Reviews* 2011;40:3656-3676.
239. Noether E; Tavel M (translator). *Invariant Variation Problems. Transport Theory and Statistical Physics* 1971;1:186-207.
240. Sokolov VI. Concept of a Stereochemical System. *Russian Journal of Organic Chemistry* 2002;38:1553-1563.
241. István Hargittai I. *Symmetry: Unifying Human Understanding*. Pergamon Press 1986.
242. Tverdislov VA, Yakovenko LV and Zhavoronko AA, Chirality as a problem of biochemical physics. *Russian Journal of General Chemistry* 2007;77:1994-2005.
243. Tverdislov VA. Chirality as an Instrument of Stratification of Hierarchical Systems in Animate and Inanimate Nature. Cornell University Library 2012.
244. Kimura T, Hamase K, Miyoshi Y, et al. Chiral amino acid metabolomic for novel biomarker screening in the prognosis of chronic kidney disease. *Scientific Reports* 2016;6:26137.
245. Gunturkun O and Ocklenburg S. Ontogenesis of Lateralization. *Neuron* 2017;94:249-263.
246. Celeman PD and Mastroeni D. A Call for new approaches to Alzheimer's disease research. *Neuro-Biology of Aging* 2017;57:iii-iv.
247. Rothschild G and Mizrahi A. Global Order and Local Disorder in Brain Maps. *Annual Review of Neuroscience* 2015;38:247-268.
248. Niell CM. Cell Types, Circuits, and Receptive Fields in the Mouse Visual Cortex. *Annual Review of Neuroscience* 2015;38:413-431.
249. Seabrook TA, Burbridge TJ, Crair MC and Huberman AD. Architecture, Function, and Assembly of the Mouse Visual System. *Annual Review of Neuroscience* 2017;40:499-538.
250. Kaplan G. Audition and Hemispheric Specialization in Songbirds and New Evidence from Australian Magpies. *Symmetry* 2017;9:99.
251. Schmitz J, Lor S, Klose R, Güntürkün O and Ocklenburg S. The Functional Genetics of Handedness and Language Lateralization: Insights from Gene Ontology, Pathway and Disease Association Analyses. *Frontiers in Psychology* 2017;8:1144
252. Ocklenburg S, Schmitz J, Moinfar Z, et al. Epigenetic regulation of lateralized fetal spinal gene expression underlies hemispheric asymmetries. *eLIFE* 2017;6:e22784.
253. Towse CL, Hopping G, Vulovic I and Daggett V. Nature versus design: the conformational propensities of D-amino acids and the importance of side chain chirality. *Protein Engineering, Design and Selection* 2014;27:447-455.